



Original communication

Determination of ethyl glucuronide in human hair by hydrophilic interaction liquid chromatography–tandem mass spectrometry[☆]

Fadile Yaldiz, MSc, Specialist^a, Nebile Daglioglu, PhD, Forensic Toxicologist^{b,*},
Ahmet Hilal, MD Assoc. Professor, Lecturer^b, Alper Keten, M.D, Medical Examiner^c,
Mete Korkut Gülmen, M.D Professor, Lecturer^b

^a Mersin University, Advanced Technology Education, Research and Application Center, Mersin, Turkey

^b Cukurova University, School of Medicine, Department of Forensic Medicine, Adana, Turkey

^c Department of Forensic Medicine, Ataturk Education and Research Hospital, Ankara, Turkey

ARTICLE INFO

Article history:

Received 3 September 2012

Received in revised form

10 June 2013

Accepted 27 June 2013

Available online 7 August 2013

Keywords:

Ethyl Glucuronide

Hair

Alcohol addiction

HILIC

ABSTRACT

Ethyl glucuronide (EtG) is a direct metabolite of ethanol and has been utilized as a marker for alcohol intake. This study presents development, validation and application of a new hydrophilic interaction liquid chromatography–tandem mass spectrometry (HILIC–MS/MS) method for the analysis of EtG in human hair samples. The linearity was assessed in the range of 5–2000 pg/mg hair, with a correlation coefficient of >0.99. The method was selective and sensitive, with a limit of detection (LOD) and limit of quantitation (LOQ) of 0.05 pg/mg and 0.18 pg/mg in hair, respectively. Differently from the extraction procedures in the literature, a fast and simple liquid–liquid method was used and highest recoveries and cleanest extracts were obtained.

The method was successfully applied to 30 human hair samples which were taken from those who state they consume alcohol. EtG concentrations in the hair samples of alcohol users participated in this study, ranged between 1.34 and 82.73 pg/mg. From the concentration of EtG in hair strands 20 of the 30 subjects can be considered regular moderate drinkers.

© 2013 Elsevier Ltd and Faculty of Forensic and Legal Medicine. All rights reserved.

1. Introduction

Overconsumption of alcohol is a contributing factor medico-legal case.¹ It is a fact that alcohol is a contributing factor for traffic accidents, serious traumas, natural deaths and the other medico-legal cases. Furthermore, alcoholism is one of the most frequent addictions. In such cases, the most frequently used samples for ethanol analysis are blood, urine, saliva, and breath. Additionally, compared to urine or blood analysis, hair analysis offers a much larger window of detection for alcohol consumption. Ethyl glucuronide (EtG) is a product of the non-oxidative ethanol metabolism. EtG is a nonvolatile, water-soluble, direct metabolite of ethanol, which can be detected in body fluids, tissues and hair. Hair analysis has become an important diagnostic tool for the determination of chronic alcohol consumption since EtG can accumulate

in hair. Its first identification was reported by Sachs et al., in 1993.² Several methods to determine EtG concentration in human hair have been recently developed.^{3–7}

Previously conducted studies have utilized analytical methods based on liquid chromatography (LC) with mass spectrometry (MS).⁸ LC–MS/MS methods do not need derivatizations, and are thus less complex and time consuming. Recently, the use of hydrophilic interaction liquid chromatography (HILIC) coupled to MS, developed for the separation of polar compound. HILIC was applied successfully for the analysis of EtG in hair.^{9,10}

The aims of this study were to determine EtG in human hair samples using HILIC, and to find out the correlation between alcohol intake and EtG levels in hair samples.

2. Materials and methods

2.1. Hair samples

Thirty hair samples included in the study belonged to males aged between 25 and 63 years with a mean age of 41.77 and median age of 42.50 years. A written informed consent to participate in the study was obtained from all the individuals. Before collecting the

[☆] This study was presented in “22nd Congress of the International Academy of Legal Medicine, İstanbul, Türkiye, July 5–8, 2012” as an oral presentation.

* Corresponding author. Tel.: +90 530 886 30 94; fax: +90 322 338 63 06.

E-mail addresses: fadileyener@gmail.com (F. Yaldiz), nebile@hotmail.com (N. Daglioglu), ahilal@cu.edu.tr (A. Hilal), alperketen@gmail.com (A. Keten), mete.gulmen@gmail.com (M.K. Gülmen).

Table 1
Matrix effect.

Analyte	Theoretical concentration (ng/ml)	Matrix effect (%)	Relative matrix effect RSD (%)
EtG	20	105	3.5
	500	100	2.7

hair samples, each participant was requested to complete an anonymous questionnaire. This included a series of questions such as daily intake of ethanol, age, and sex. The study was approved by the Ethics Committee of Cukurova University. Hair specimens were collected by fixing a strand of hair on the posterior region of vertex and by cutting it as close to the skin as possible. Blank hair was obtained from young children free from alcohol. The samples were stored in aluminum foil until analyses.

2.2. Chemicals and reagents

EtG and d_5 -EtG were purchased from Medichem (Germany). Methanol and acetonitrile were obtained from Merck (Darmstadt, Germany). All chemicals were of analytical grade or highest purity available.

Stock solutions of EtG (1 mg/ml) and d_5 -EtG (0.5 mg/ml) were prepared in methanol and they were stored at $-20\text{ }^{\circ}\text{C}$. Working standard solutions used for calibration and quality control samples were prepared by 0.015, 0.025, 0.05, 0.1, 0.2, 0.5, 1, 5 and $10\text{ }\mu\text{g/ml}$. All working solutions were stored in a refrigerator when not in use.

2.3. Sample preparation

Hair samples were washed twice using water (5 ml). Each strand was cut into small pieces ($>1\text{ mm}$) by scissors and 200 mg of the hair was weighted in 10-ml conical-bottom glass tubes and spiked with $50\text{ }\mu\text{l}$ internal standard solution (d_5 -EtG, 2.5 mg/l in methanol). The samples were vortexed for 1 min and samples were ultrasonicated for 15 min in 2 ml acetonitrile/water solution (80:20), centrifuged (3500 rpm for 5 min). Next, the supernatant was filtrated by nylon filter and injected into the LC–MS/MS.

2.4. Instrumentation

Chromatographic separation was carried out with an LC system comprised of an Agilent 1200 series binary pump, vacuum degasser and auto-sampler (Agilent, Germany). The analyte and internal standard were eluted on a Zorbax Hilic Plus, Rapid Resolution ($50\text{ mm} \times 4.6\text{ mm}$, $3.5\text{ }\mu\text{m}$, Agilent, USA). Mobile phase composed by (A) ammonium acetate 1 mM in water and (B) acetonitrile. The

separation was achieved by isocratic elution. The injection volume $10\text{ }\mu\text{l}$ was used in all cases.

Analyses were performed on an Agilent (USA) triple quadrupole mass spectrometer model 6460, equipped with an atmospheric pressure pneumatically assisted electrospray ionization source. The interface was operated in negative ionization mode. High purity nitrogen served both as collision-induced dissociation gas, drying gas (set at 11 L/min) and nebulizer gas (set at 50 psi). The source temperature was maintained at $350\text{ }^{\circ}\text{C}$, the capillary voltage was set at 4000 V and the fragment or voltage at 135 V after optimization by injecting EtG standard. Analysis was performed by multi-reaction monitoring, using $221 \rightarrow 75$, $221 \rightarrow 85\text{ amu}$ for EtG and $226 \rightarrow 75$, $226 \rightarrow 85\text{ amu}$ for d_5 -EtG.

2.5. Validation

The method was validated for selectivity, linearity and sensitivity, precision and accuracy and for recovery.

Seven-point calibration curves were prepared by spiking the weighted amount of blank hair with standard solution to obtain the concentration range of 0–5–20–50–100–500–1000–2000 pg/mg for EtG. To assess linearity, a calibration curve was prepared in replicate ($n = 3$) and analyzed. The acceptance criteria were: a correlation coefficient (r) and determination coefficient (r^2) > 0.99 , and a precision and accuracy, for the back calculated concentrations of the calibration points, within $\pm 15\%$.

The limit of detection (LOD) was 0.05 pg/mg, with an S/N ratio of 3. The limit of quantitation (LOQ) was 0.18 pg/mg. Under the chromatographic conditions used, there was no interference with the analytes by chemicals or any extractable endogenous materials present in hair.

The extraction recovery tests were performed by analyzing extracted hair samples and diluted standard solutions at equal concentrations. Five replications at low and high concentrations were prepared. The recoveries at 20 and 1000 pg/mg were 97.84% and 100%, respectively.

The matrix effect was evaluated according to the following procedure¹¹: the analyte signal in the spiked water was compared with the analyte signal in the matrix fortified after extraction and the matrix effect percentage was calculated as $\% = (\text{extracted matrix height/water height}) \times 100$. Five replicates of hair extracts were analyzed (Table 1).

3. Results and discussion

The chromatograms of blank hair sample and of a blank hair sample spiked with EtG (20 pg/mg) concentration, extracted as described above and separated in HILIC conditions, are presented in Fig. 1.




Fig. 1. Chromatograms of blank hair sample and a blank spiked with EtG at 20 pg/mg.




Fig. 2. Chromatogram of a real sample (EtG: 82.73 pg/mg).

A total of 30 human hair samples were analyzed for EtG. The EtG concentrations were between 1.34 and 82.73 pg/mg (mean 13.5 ± 15.25 pg/mg, median 9.5 ± 15.25 pg/mg). Fig. 2 shows the chromatogram obtained after extraction of a hair from an alcohol user.

EtG, a metabolite of ethanol, is a valuable marker for alcohol intake; thus, it is of high importance in forensic and clinical cases involving alcohol. Skipper et al.¹² point out that a prerequisite for a reliable test is the ability to show the use of alcohol at least in a few days after its consumption. According to a report by WHO, men who take at least 40 g of alcohol per day and women who use at least 20 g of alcohol are accepted as heavy drinkers.¹³

The aim of the hair test is to reveal the suspicious alcohol use and determine the consumption of alcohol by the people who are previously known to be alcohol users. The difference between social drinkers and alcohol addicts is specified on the basis of EtG cut-off values. In order to avoid different interpretations done by different laboratories, the Society of Hair Analysis (2012) reported the value of EtG cut-off as 7 pg/mg.¹⁴ The concentrations above this value show repeated alcohol consumption. In our study, only twelve samples showed concentrations above 7 pg/mg. The other values were found to be below the cut-off value. This clearly shows that those people consumed alcohol repeatedly.

Pragst et al.¹⁵ detected EtG in hair samples obtained from three heavy drinkers. Similarly, Alt et al.¹⁶ identified EtG concentrations ranging between 218 and 4025 pg/mg, in 16 postmortem hair samples of alcohol addicts. They also found that EtG concentrations ranged between 119 and 388 pg/mg in the patients under treatment. However, they found no evidence of EtG in the hair samples of children and of social drinkers who consume alcohol below 20 g per day.

The mean value of EtG concentrations in the currently studied hair samples was 13.5 ± 15.25 pg/mg. The common characteristics

of these samples were that these people were social drinkers. The differences in the concentrations of EtG in hair samples was related to daily consumption of alcohol as shown by Spearman correlation coefficient ($r = 0.668$, $p < 0.001$) as shown in Fig. 3. These findings were compatible with those of by Politi et al.,¹³ Appenzeller et al.⁷ and Kronstrand et al.¹⁷; however, no correlation between the amount of consumed alcohol and the EtG concentrations was observed in hair samples, in the study by Janda et al.¹⁸ One of our study limitations was participant's hair color because some studies^{7,19} have also shown that hair pigmentation does not influence incorporation of EtG in hair therefore hair color was not included in this study.




Fig. 3. Relationship between EtG concentration and ethanol daily intake.

In the present study, we demonstrated the successful determination of EtG levels by LC/MS/MS using a HILIC column. However, for EtG, which is an extremely polar compound, the use of hydrophilic column was thought to be preferable. Through HILIC column use, the detection of a very low level of EtG could be achieved (LOQ: 0.18 pg/mg) in hair samples.

4. Conclusion

In this study, a selective and sensitive identification and quantification method for EtG in human hair samples was developed using HILIC/MS/MS. The method was applied to real samples and was able to rapidly quantify EtG also in the hair samples of alcohol users.

Ethical approval

None.

Funding

Cukurova University Research Foundation provided the financial support for this work. (Grant no: TF 2009YL15).

Conflict of interest

None.

Competing interests

There are no competing interests for any of authors.

Contributorship

All authors included in the authors list have contributed to the data collection, data evaluation, writing of manuscript, language use etc. And no professionals other than authors contributed to the any process of during article preparation.

References

1. WHO.org [internet]. Geneva: *Global status report on alcohol and health*. [Updated 2011 February 11]. Available from: <http://www.who.int/>.
2. Sachs H. History of hair analysis. *Forensic Sci Int* 1997;**84**:7–16.
3. Xu RN, Fan L, Rieser MJ, Tawakol A. Recent advances in high-throughput quantitative bioanalysis by LC–MS/MS. *J Pharm Biomed Anal* 2007;**44**:342–55.
4. Yegles M, Labarthe A, Auwarter V, Hartwig S, Vater H, Wennig R, et al. Comparison of ethyl glucuronide and fatty acid ester concentrations in hair of alcoholics, social drinkers and teetotalers. *Forensic Sci Int* 2004;**145**:167–73.
5. Auwarter V, Sporkert F, Hartwig S, Pragst F, Vater F. Fatty acid ethyl esters in hair as markers of alcohol consumption, segmental hair analysis of alcoholics, social drinkers, and teetotaler. *Clin Chem* 2001;**47**:2114–23.
6. Hoiseith G, Morini L, Poletini A. Ethyl glucuronide in hair compared with traditional alcohol biomarkers – a pilot study of heavy drinkers referred to an alcohol detoxification unit. *Alcohol Clin Exp Res* 2009;**33**:812–6.
7. Appenzeller BMR, Schuman M, Yegles M, Wennig R. Ethyl glucuronide concentration in hair is not influenced by pigmentation. *Alcohol Alcohol* 2007;**42**:326–7.
8. Keten A, Tümer AR, Balseven-Odabaşı A. Measurement of ethyl glucuronide in vitreous humor with liquid chromatography-mass spectrometry. *Forensic Sci Int* 2009;**193**:101–5.
9. Kintz P, Villain M, Cirimele V. Hair analysis for drug detection. *Ther Drug Monit* 2006;**28**:442–6.
10. Tarcomnicu I, Van Nuijs ALN, Aerts K, De Doncker M, Covaci A, Neels H. Ethyl glucuronide determination in meconium and hair by hydrophilic interaction liquid chromatography-tandem mass spectrometry. *Forensic Sci Int* 2010;**196**:121–7.
11. Matuszewski BK, Constanzer ML, Chavez-Eng CM. Strategies for the assessment of matrix effect in quantitative bioanalytical methods based on HPLC-MS/MS. *Anal Chem* 2003;**75**:3019–30.
12. Skipper GE, Weinman W, Wurst FM. Ethylglucuronide (EtG): a new marker to detect alcohol use in recovering physicians. *J Med Licensure Discipline* 2004;**90**:14–7.
13. Politi L, Morini L, Poletini A. Ethyl glucuronide in Hair: is it a reliable marker of chronic high levels of alcohol consumption? *Methods Tech* 2006;**101**:1408–12.
14. Soht.org [internet]. *Use of Alcohol Markers in Hair for Abstinence Assessment 2012*, [updated 2012 February 18]. Available from: www.soht.org.
15. Pragst F, Spiegel K, Sporkert F, Bohnenkamp M. Are there possibilities for the detection of chronically elevated alcohol consumption by hair analysis? A report about the state of investigation. *Forensic Sci Int* 2000;**107**:201–23.
16. Alt A, Janda I, Seidl S, Wurst FM. Determination of ethyl glucuronide in hair samples. *Alcohol Alcohol* 2000;**35**:313–4.
17. Kronstrand R, Brinkhagen L, Nyström FH. Ethyl glucuronide in human hair after daily consumption of 16 or 32 g of ethanol for 3 months. *Forensic Sci Int* 2012;**215**:51–5.
18. Janda L, Weinmann W, Kuehnle T, Lahode M, Alt A. Determination of ethyl glucuronide in human hair by SPE and LC-MS/MS. *Forensic Sci Int* 2002;**128**:59–65.
19. Kharbouche H, Steiner N, Morelato M, Staub C, Boutrel B, Mangin P, et al. Influence of ethanol dose and pigmentation on the incorporation of ethyl glucuronide into rat hair. *Alcohol* 2010;**44**:507–14.